

Cardiovascular Effects of 2-Arachidonoyl Glycerol in Anesthetized Mice

Zoltán Járαι, Jens A. Wagner, Sravan Kumar Goparaju, Lei Wang, Raj K. Razdan, Takayuki Sugiura, Ann M. Zimmer, Tom I. Bonner, Andreas Zimmer, George Kunos

Abstract—Cannabinoids, including the endogenous ligand anandamide, elicit pronounced hypotension and bradycardia through the activation of CB1 cannabinoid receptors. A second endogenous cannabinoid, 2-arachidonoyl glycerol (2-AG), has been proposed to be the natural ligand of CB1 receptors. In the present study, we examined the effects of 2-AG on mean arterial pressure and heart rate in anesthetized mice and assessed the role of CB1 receptors through the use of selective cannabinoid receptor antagonists and CB1 receptor knockout (CB1^{-/-}) mice. In control ICR mice, intravenous injections of 2-AG or its isomer 1-AG elicit dose-dependent hypotension and moderate tachycardia that are unaffected by the CB1 receptor antagonist SR141716A. The same dose of SR141716A (6 nmol/g IV) completely blocks the hypotensive effect and attenuates the bradycardic effect of anandamide. 2-AG elicits a similar hypotensive effect, resistant to blockade by either SR141716A or the CB2 antagonist SR144528, in both CB1^{-/-} mice and their homozygous (CB1^{+/+}) control littermates. In ICR mice, arachidonic acid (AA, 15 nmol/g IV) elicits hypotension and tachycardia, and indomethacin (14 nmol/g IV) inhibits the hypotensive effect of both AA and 2-AG. Synthetic 2-AG incubated with mouse blood is rapidly (<2 minutes) and completely degraded with the parallel appearance of AA, whereas anandamide is stable under the same conditions. A metabolically stable ether analogue of 2-AG causes prolonged hypotension and bradycardia in ICR mice, and both effects are completely blocked by SR141716A, whereas the same dose of 2-AG-ether does not influence blood pressure and heart rate in CB1^{-/-} mice. These findings are interpreted to indicate that exogenous 2-AG is rapidly degraded in mouse blood, probably by a lipase, which masks its ability to interact with CB1 receptors. Although the observed cardiovascular effects of 2-AG probably are produced by an arachidonate metabolite through a noncannabinoid mechanism, the CB1 receptor-mediated cardiovascular effects of a stable analogue of 2-AG leaves open the possibility that endogenous 2-AG may elicit cardiovascular effects through CB1 receptors. (*Hypertension*. 2000;35:679-684.)

Key Words: receptors ■ mice ■ hypotension ■ cannabinoid

It has long been known that cannabinoids have powerful cardiovascular effects. In anesthetized rats, Δ^9 -tetrahydrocannabinol (THC) causes prolonged hypotension and bradycardia.¹ In 1992, arachidonyl ethanolamide (anandamide) extracted from porcine brain was identified as an endogenous cannabinoid based on its ability to bind to the brain cannabinoid receptor and to mimic the neurobehavioral effects of THC.² Anandamide also elicits hypotension and bradycardia in anesthetized rats through a peripheral mechanism.³⁻⁵ These effects are mediated by CB1 cannabinoid receptors,⁶ as indicated by their susceptibility to blockade by SR141716A,^{3,5} a selective CB1 receptor antagonist,⁷ and by their absence in CB1 receptor knockout (CB1^{-/-}) mice.⁸ In 1995, 2-arachidonoyl glycerol (2-AG), found in canine gut⁹ and rat brain,¹⁰ was identified as another endogenous canna-

binoid. The concentration of 2-AG in the brain is several hundredfold higher than that of anandamide,^{10,11} and, unlike anandamide, which often acts as a partial agonist,¹²⁻¹⁵ 2-AG is a full agonist at CB1 receptors.^{11,16,17} This has led to the proposal that the CB1 receptor is a 2-AG receptor.¹⁷

Endogenous cannabinoids also have been implicated in the local control of vascular tone. In certain vascular preparations, SR141716A inhibits the vasodilation induced by the endothelium-derived hyperpolarizing factor,^{18,19} and anandamide has been identified in vascular endothelial cells.²⁰ SR141716A also antagonizes hemorrhagic²¹ and endotoxin-induced hypotension,²² and circulating macrophages produce anandamide under such conditions, which suggests a paracrine mechanism in the control of vascular tone.^{21,22} Recently, 2-AG also has been identified in vascular endothelial

Received June 30, 1999; first decision August 26, 1999; revision accepted September 27, 1999.

From the Department of Pharmacology and Toxicology, Medical College of Virginia of Virginia Commonwealth University, Richmond (Z.J., J.A.W., S.K.G., L.W., G.K.); Organix, Inc, Woburn, Mass (R.K.R.); the Faculty of Pharmaceutical Sciences, Teikyo University, Sagamiko, Kanagawa, Japan (T.S.); and the Section on Genetics, National Institute of Mental Health, Bethesda, Md (A.M.Z., T.I.B., A.Z.). J.A.W. is now with the Department of Medicine, University of Wuerzburg, Wuerzburg, Germany.

Correspondence to Dr George Kunos, Department of Pharmacology and Toxicology, MCV/VCU, 410 North 12th St, PO Box 980613, Richmond, VA 23298. E-mail gkunos@hsc.vcu.edu

© 2000 American Heart Association, Inc.

Hypertension is available at <http://www.hypertensionaha.org>




Figure 1. Effects of 2-AG (A) and anandamide (B) on mean BP and HR in ICR mice in the absence or presence of SR141716A. 2-AG (26.4 nmol/g) or anandamide (11.5 nmol/g) was injected intravenously at 0 minutes, 10 minutes after the intravenous injection of vehicle (○) or 6 nmol/g SR141716A (●). Points and bars represent mean \pm SE, $n=5$ to 19. *Significant difference from baseline.

cells,^{23,24} and in platelets the concentration of 2-AG was found to increase in response to endotoxin treatment.²² Thus 2-AG also may play a role in the local control of vascular tone by the activation of SR141716A-sensitive receptors.

We²² and others²⁴ have reported that intravenous injection of 2-AG in anesthetized rats causes dose-dependent hypotension. However, this effect was antagonized by SR141716A less effectively than anandamide-induced hypotension,²² and, unlike anandamide, which causes CB1 receptor-mediated bradycardia,⁵ 2-AG elicits tachycardia unaffected by SR141716A.²² In the present study we analyzed the cardiovascular effects of 2-AG in anesthetized mice and assessed the role of CB1 receptors through the use of selective antagonists and CB1^{-/-} mice. The results indicate that 2-AG is rapidly degraded in mouse blood, and the observed hypotensive effect does not involve CB1 receptors but probably is caused by an arachidonic acid (AA) metabolite. In contrast, a metabolically stable analogue of 2-AG causes CB1 receptor-mediated hypotension and bradycardia.

Methods

Cardiovascular Measurements

Ninety male ICR mice (25 to 30 g) from Harlan (Indianapolis, Ind) were used. Male and female CB1^{-/-} mice and their homozygous CB1^{+/+} littermates (64 animals) have been described.²⁵ Animals were kept under a 12:12 hours light/dark cycle, fed standard mouse chow, and had access to drinking water ad libitum. Mice were anesthetized with 60 mg/kg IP sodium pentobarbital, supplemented as needed to maintain stable anesthesia. The jugular vein and the carotid artery were cannulated with PE-10 tubing filled with heparinized saline for intravenous drug injections and monitoring blood pressure (BP) and heart rate (HR), respectively. The arterial cannula was connected to a pressure transducer and physiograph. HR was derived from the pressure pulse with the use of a tachograph preamplifier. The procedures used were approved by the institutional animal care and usage committee.

Analysis of 2-AG and Anandamide by Liquid Chromatography/Mass Spectrometry

For assessing the stability of 2-AG and anandamide in mouse blood, 200- μ L aliquots of heparinized blood were withdrawn from ICR

mice and incubated with 100 μ g of synthetic 2-AG (final concentration 1.32 mmol/L) or 100 μ g of anandamide (1.44 mmol/L) for 1 to 5 minutes at 37°C. At the end of the incubation, the whole blood was extracted 3 times with 500 μ L of cold diethyl ether + 20 μ L 1.0N HCl; the organic phase was dried under a stream of nitrogen and resuspended in 100 μ L methanol. The sample was fractionated by reversed-phase high-performance liquid chromatography on an ODS column (Supelcosil, 5 μ m, 4.6 mm \times 15 cm), with the use of a mobile phase of methanol/water/acetic acid (85:15:0.03 vol/vol/vol) at a flow rate of 1 mL/min on a Waters 2690 system. This was followed in line by mass spectroscopic analysis on a Micromass Quattro II mass spectrometer equipped with an atmospheric pressure chemical ionization source. Two selected ions were simultaneously monitored (alternating at 0.2 seconds), as specified in the legend of Figure 5.

Drugs

2-AG and 1- (or 3-)-arachidonoyl glycerol (1-AG) were synthesized as described earlier.²⁶ The synthesis of an ether-linked analogue of 2-AG, 2-eicosatetraenylglycerol (2-AG-ether), has been detailed elsewhere.¹⁷ The structure of all 3 compounds was confirmed by nuclear magnetic resonance and electron impact mass spectrometry. SR141716A⁷ (*N*-[piperidin-1-yl]-5-[4-chlorophenyl]-1-[2,4-dichlorophenyl]-4-methyl-1H-pyrazol-3-carboxamide HCl) and the CB2 receptor antagonist SR144528²⁷ (*N*-[(1*S*)-endo-1,3,3-trimethyl bicyclo[2.2.1]heptan-2-yl]-5-(4-chloro-3-methylphenyl)-1-(4-methyl-benzyl)-pyrazole-3-carboxamide) were gifts from Sanofi Recherche (Montpellier, France). Anandamide was provided by Dr Billy R. Martin.

Statistical Analyses

The dose- or time-dependent effects of drugs on BP and HR and on neurobehavioral parameters were analyzed by ANOVA followed by Tukey's post hoc test. Differences from corresponding baseline values with a value of $P < 0.05$ were considered statistically significant.

Results

Bolus intravenous injection of 26.4 nmol (10 μ g)/g 2-AG in ICR mice caused hypotension and moderate tachycardia that lasted \approx 10 minutes. Unexpectedly, neither effect was influenced by pretreatment with 6 nmol (3 μ g)/g SR141716A (Figure 1A). The same dose of SR141716A nearly completely blocked the hypotension and attenuated the bradycar-




Figure 2. Dose-dependence and lack of inhibition by SR141716A of the hypotensive effect of 2-AG and 1-AG in ICR mice. Columns and bars represent mean \pm SE (black: controls, $n=6$ to 15; shaded: after 6 nmol/g SR141716A, $n=4$ to 9). *Significant difference from baseline.

dia elicited by an equihypotensive dose (11.5 nmol [$4 \mu\text{g}$]/g) of anandamide (Figure 1B). Basal BP in the absence or presence of SR141716A was 74 ± 3 versus 76 ± 5 mm Hg, and HR was 365 ± 13 versus 414 ± 16 bpm. The dose dependence of the hypotensive response to 2-AG is illustrated in Figure 2A. At none of the doses tested was any inhibition by SR141716A observed. 1-AG, which has been described to have similar potency as 2-AG in producing cannabinoid effects,¹¹ also caused dose-dependent hypotension, which was also not blocked by SR141716A pretreatment (Figure 2B). In fact, SR141716A potentiated the effect of 1-AG, which was significant at the 2.6 nmol/g dose.

Because the lack of inhibition by SR141716A called into question the role of CB1 receptors in the effects of 2-AG, we tested these effects in $\text{CB1}^{-/-}$ and $\text{CB1}^{+/+}$ mice (Figure 3). In both types of animals, 26.4 nmol/g of 2-AG caused hypotension and tachycardia similar to the effects seen in ICR mice, although the hypotension was somewhat less in $\text{CB1}^{-/-}$ than in $\text{CB1}^{+/+}$ animals. Nevertheless, the effect of SR141716A (6 nmol/g IV) on the hypotensive effect of 2-AG was similar in the 2 groups: no inhibition was observed. In $\text{CB1}^{+/+}$ mice, basal BP in the absence or presence of SR141716A was 81 ± 6 versus 84 ± 9 mm Hg and basal HR was 278 ± 19 versus 416 ± 10 bpm. In $\text{CB1}^{-/-}$ mice, the corresponding values were 75 ± 4 versus 75 ± 1 mm Hg and 331 ± 20 versus 413 ± 28

bpm. Although the tachycardic effect of 2-AG was reduced by SR141716A in both $\text{CB1}^{-/-}$ and $\text{CB1}^{+/+}$ mice, this may have been related to the marked increase in basal HR caused by SR141716A in both groups. Pretreatment of CB1 receptor knockout mice with the CB2 receptor antagonist SR144528 (10.5 nmol [$5 \mu\text{g}$]/g IV, Figure 3), which did not affect basal BP or HR, also failed to affect the responses to 2-AG.

In contrast to SR141716A, indomethacin (14 nmol [$5 \mu\text{g}$]/g IV) pretreatment of ICR mice significantly inhibited the hypotensive response to 2-AG, whereas the moderate tachycardia was unaffected (Figure 4A). AA (15 nmol [$5 \mu\text{g}$]/g IV) also elicited hypotension and moderate tachycardia, and the hypotension but not the tachycardia was inhibited by 14 nmol/g indomethacin (Figure 4B). On the other hand, SR141716A pretreatment (6 nmol/g IV) failed to influence the effects of AA (Figure 4B), and, conversely, indomethacin (14 nmol/g IV) did not influence the hypotensive response to 11.5 nmol/g anandamide (18 ± 2 vs 19 ± 3 mm Hg, $n=5$, before and after indomethacin, respectively). Because this suggested that 2-AG may break down to release AA, which could be further metabolized into a hypotensive cyclooxygenase product(s), we tested the stability of 2-AG in mouse blood. Authentic 2-AG or anandamide (100 μg each) was added to 0.2-mL aliquots of heparinized mouse blood at 37°C , the blood samples were extracted after




Figure 3. Effects of 2-AG on mean BP (left) and HR (right) in $\text{CB1}^{-/-}$ (top) and $\text{CB1}^{+/+}$ mice (bottom). 2-AG (26.4 nmol/g) was injected intravenously before (\circ) and 10 minutes after the intravenous injection of 6 nmol/g SR141716A (\bullet) or 10.4 nmol/g SR144528 (\blacksquare). Points and bars represent mean \pm SE, $n=6$ to 9. *Significant difference from baseline.




Figure 4. Effect of indomethacin on 2-AG-induced (A) and AA-induced (B) changes in mean BP and HR in ICR mice. 2-AG (26.4 nmol/g) or AA (15 nmol/g) was injected intravenously 10 minutes after the intravenous injection of vehicle (○) or 14 nmol/g indomethacin (▲). AA was also tested after pretreatment with 6 nmol/g SR141716A (●). Points and bars represent mean \pm SE, $n=5$ to 6. *Significant difference ($P < 0.05$) from baseline.

various time intervals with diethyl ether, and the extracts were analyzed by liquid chromatography/mass spectrometry. As illustrated in the left panels of Figure 5, 2-AG disappeared from the blood within 2 minutes, with the parallel appearance of AA. In contrast, most of the anandamide added could be recovered unmetabolized after similar incubations (Figure 5, right panels) and even after a 5-minute incubation (not shown). These findings confirm the extreme instability of 2-AG in mouse blood, whereas anandamide appears to be more stable under similar conditions.

Because of its instability, 2-AG may not reach its site of action in the vasculature in amounts sufficient to activate CB1 receptors. Therefore we analyzed the cardiovascular effects of a metabolically stable ether analogue of 2-AG^{17,24} in ICR mice and in CB1^{-/-} mice. As seen in Figure 6A,

intravenous injection into ICR mice of 56 nmol (20 μ g)/g 2-AG-ether elicited prolonged hypotension and bradycardia, and both effects were completely blocked by pretreatment with 6 nmol/g SR141716A. Similar effects were observed in CB1^{+/+} mice (data not shown), whereas 2-AG-ether failed to elicit any change in BP and caused a modest increase in HR in CB1^{-/-} mice (Figure 6B). This further confirms the role of CB1 receptors in the effects of 2-AG-ether.

Discussion

The results presented in this article illustrate an unexpected difficulty in documenting the role of cannabinoid receptors in a biological effect of 2-AG on its intravenous administration to mice. Cannabinoids, including anandamide, evoke hypotension and bradycardia by activating CB1 receptors.^{3,5} Al-




Figure 5. LC/MS analysis of 2-AG and anandamide incubated with mouse blood. Two selected ions were monitored simultaneously: m/z 379.4 for 2-AG ions and m/z 305.4 for AA ions in the left panels and m/z 348.4 for anandamide ions and m/z 305.4 for AA ions in the right panels. Samples used were extracts of 200 μ L drug-free whole mouse blood (A); of 100 μ g authentic 2-AG or anandamide dissolved in 200 μ L buffer (B); and of 200 μ L blood samples incubated with 100 μ g 2-AG or anandamide for 1 minute (C) or 2 minutes (D).




Figure 6. Effects of 2-AG-ether on BP and HR in ICR mice (A) and in CB1^{-/-} mice (B). 2-AG-ether (56 nmol/g) was injected intravenously at 10 minutes after the injection of vehicle (○) or, in ICR mice only, 10 minutes after the intravenous injection of 6 nmol/g SR141716A (●). Points and bars represent mean \pm SE, $n=4$ to 6. *Significant difference from baseline.

though 2-AG also elicits hypotension, it increases rather than decreases HR, which is unusual in view of its documented ability to interact with CB1 receptors. The inability of SR141716A to inhibit the hypotensive effect of 2-AG and the persistence of such an effect in CB1^{-/-} mice clearly indicate the lack of CB1 receptor involvement. 2-AG induces similar tachycardia in CB1^{+/+} and CB1^{-/-} mice, which discounts the involvement of CB1 receptors in this effect as well. Although the tachycardic effect of 2-AG was reduced by SR141716A in both groups of mice, the apparent inhibition probably was related to the substantial increase in basal HR after SR141716A treatment. SR141716A does not cause tachycardia in rats,³ and the presence of the tachycardic effect in both CB1^{-/-} and CB1^{+/+} mice rules out the involvement of CB1 receptors. The mechanism of SR141716A-induced tachycardia in mice thus remains unclear.

We present 3 lines of evidence to indicate that these paradoxical findings are due to the rapid degradation of 2-AG in mouse blood and the subsequent generation of a hypotensive arachidonate metabolite whose actions are not mediated by cannabinoid receptors. First, the effects of 2-AG could be mimicked by AA, and the effects of both 2-AG and AA were antagonized by indomethacin. Because indomethacin did not influence the hypotensive response to anandamide, its effect cannot be attributed to a nonspecific interaction with CB1 receptors but is more likely to be caused by inhibition of the generation of an as-yet-unidentified hypotensive cyclooxygenase product.

The second evidence is the rapid disappearance of 2-AG from mouse blood. The parallel appearance of AA suggests that 2-AG is rapidly hydrolyzed to yield AA and glycerol. Because the breakdown is essentially complete in 2 minutes, well before the peak of the hypotensive response or a significant increase in HR develops, it is very unlikely that 2-AG would be responsible for these effects. The finding that anandamide is much more stable under the same conditions is consistent with its CB1-receptor-mediated hypotensive and bradycardic effects. The differential stability of these two

endocannabinoids also suggests that the rapid breakdown of 2-AG is not catalyzed primarily by an amidohydrolase, which can degrade both anandamide and 2-AG,²⁸ but rather by a lipase such as *sn*-2 monoacylglycerol lipase, which may be involved in the degradation of 2-AG but not of anandamide.^{29,30} Further studies may be warranted to analyze the activity of this enzyme in blood.

The third line of evidence is the effects of a metabolically stable ether analogue of 2-AG. Not only did this compound produce hypotension as well as bradycardia, as expected from an agonist of CB1 receptors,⁵ but both effects were completely blocked by a dose of SR141716A previously shown to produce selective blockade of CB1 receptors.⁷ Furthermore, both effects of 2-AG-ether were absent in CB1^{-/-} mice. In a recent study, 2-AG-ether was 2 orders of magnitude less potent and also less efficacious than 2-AG in inducing CB1 receptor-mediated calcium transients in NG108 to 15 cells.¹⁷ If a similar relation holds for the cardiovascular effects of these two compounds, 2-AG should be a highly potent hypotensive and bradycardic agent if it were protected from degradation before reaching CB1 receptors. The present findings indicate that this does not happen even with high intravenous doses of 2-AG. However, endogenous 2-AG produced and released at sites close to the receptors may escape metabolic degradation. For example, 2-AG was identified in vascular endothelial cells,²³ where its levels are increased severalfold by the muscarinic agonist carbachol,²⁴ and CB1 receptors are present in cerebrovascular smooth muscle.¹⁵ If 2-AG is produced in and released from the endothelium, it may reach CB1 receptors on smooth muscle without being degraded. CB1 receptors also have been identified in vascular endothelium,²³ and 2-AG was found in platelets²² and macrophages,³⁰ where its levels were increased 2- to 3-fold in response to bacterial lipopolysaccharide.^{22,30} Because lipopolysaccharide increases the adhesion of platelets to the endothelium, platelet-derived 2-AG could act as a "juxtacrine" regulator of vascular tone, where its proximity to an endothelial site of action could protect it from

degradation. 2-AG is highly lipid soluble, and it may remain associated with the cell membrane after its release, which could further protect it against rapid degradation by blood-borne lipases. In various tissues, 2-AG is accompanied by certain 2-acyl glycerol esters, which have no cannabinoid activity of their own but can potentiate the binding of 2-AG to CB1 receptors.³¹ This "entourage effect" has been attributed, at least in part, to the protection of 2-AG against enzymatic breakdown.³¹ Clearly, further studies are needed to explore the role of endogenous 2-AG as a cardiovascular regulator.

Acknowledgments

This work was supported by National Institutes of Health grant HL-59257 (to G.K.). Fellowship support (to Z.J.) was from Sanofi Recherche (Montpellier, France) and from the Deutsche Forschungsgemeinschaft (to J.A.W.). We thank D. Troy Bridgen for assistance with the LC/MS analyses.

References

- Vollmer RR, Cavero I, Ertel RJ, Solomon TA, Buckley JP. Role of the central autonomic nervous system in the hypotension and bradycardia induced by (-)- Δ^9 -trans-tetrahydrocannabinol. *J Pharm Pharmacol*. 1974; 26:186–198.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G, Gibson D, Mandelbaum A, Etinger A, Mechoulam R. Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science*. 1992;258:1946–1949.
- Varga K, Lake K, Martin BR, Kunos G. Novel antagonist implicates the CB₁ cannabinoid receptor in the hypotensive action of anandamide. *Eur J Pharmacol*. 1995;278:279–283.
- Varga K, Lake KD, Huangfu D, Guyenet PG, Kunos G. Mechanism of the hypotensive action of anandamide in anesthetized rats. *Hypertension*. 1996;28:682–686.
- Lake KD, Compton DR, Varga K, Martin BR, Kunos G. Cannabinoid-induced hypotension and bradycardia in rats is mediated by CB₁-like cannabinoid receptors. *J Pharmacol Exp Ther*. 1997;281:1030–1037.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*. 1990;346:561–564.
- Rinaldi-Carmona M, Barth F, Heaulme M, Shire D, Calandra B, Congy C, Martinez S, Maruani J, Neliat G, Caput D, Ferrara P, Soubrie P, Breliere JC, Le Fur G. SR141716A, a potent and selective antagonist of the brain cannabinoid receptor. *FEBS Lett*. 1994;350:240–244.
- Ledent C, Valverde O, Cossu G, Petit F, Aubert J-F, Beslot F, Böhme GA, Imperato A, Pedrazzini T, Roques BP, Vassart G, Fratta W, Parmentier M. Unresponsiveness to cannabinoids and reduced addictive effects of opiates in CB1 receptor knockout mice. *Science*. 1999;283:401–404.
- Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminski NE, Schatz AR, Gopher A, Almog S, Martin BR, Compton DR, Pertwee RG, Griffin G, Bayewitch M, Barg J, Vogel Z. Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochem Pharmacol*. 1995;50:83–90.
- Sugiura T, Kondo S, Sukagawa A, Nakane S, Shinoda A, Itoh K, Yamashita A, Waku K. 2-Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochem Biophys Res Commun*. 1995;215:89–97.
- Stella N, Schweitzer P, Piomelli D. A second endogenous cannabinoid that modulates long-term potentiation. *Nature*. 1997;388:773–778.
- Mackie K, Devane WA, Hille B. Anandamide, an endogenous cannabinoid, inhibits calcium currents as a partial agonist in N18 neuroblastoma cells. *Mol Pharmacol*. 1993;44:498–503.
- Fride E, Barg J, Levy R, Saya D, Heldman E, Mechoulam R, Vogel Z. Low doses of anandamides inhibit pharmacological effects of delta 9-tetrahydrocannabinol. *J Pharmacol Exp Ther*. 1995;272:699–707.
- Burkey TH, Quock RM, Consroe P, Ehler FJ, Hosohata Y, Roeske WR, Yamamura HI. Relative efficacies of cannabinoid CB1 receptor agonists in the mouse brain. *Eur J Pharmacol*. 1997;336:295–298.
- Gebremedhin D, Lange AR, Campbell WB, Hillard CJ, Harder DR. Cannabinoid CB1 receptor of cat cerebral arterial muscle functions to inhibit L-type Ca^{2+} channel current. *Am J Physiol*. 1999;276: H2085–H2093.
- Sugiura T, Kodaka T, Kondo S, Nakane S, Kondo H, Waku K, Ishima Y, Watanabe K, Yamamoto I. Is the cannabinoid CB1 receptor a 2-arachidonoylglycerol receptor? structural requirements for triggering a Ca^{2+} transient in NG108–15 cells. *J Biochem*. 1997;122:890–895.
- Sugiura T, Kodaka T, Nakane S, Miyashita T, Kondo S, Suhara Y, Takayama H, Waku K, Seki C, Baba N, Ishima Y. Evidence that the cannabinoid CB1 receptor is a 2-arachidonoylglycerol receptor. *J Biol Chem*. 1999;274:2794–2801.
- Randall MD, Alexander SPH, Bennett T, Boyd EA, Fry JR, Gardiner SM, Kemp PA, McCulloch AI, Kendall DA. An endogenous cannabinoid as an endothelium-derived vasorelaxant. *Biochem Biophys Res Commun*. 1996;229:114–120.
- White R, Hiley CR. A comparison of EDHF-mediated and anandamide-induced relaxations in the rat isolated mesenteric artery. *Br J Pharmacol*. 1997;122:1573–1584.
- Deutsch DG, Goligorsky MS, Schmid PC, Krebsbach RJ, Schmid HH, Das SK, Dey SK, Arreaza G, Thorup C, Stefano G, Moore LC. Production and physiological actions of anandamide in the vasculature of the rat kidney. *J Clin Invest*. 1997;100:1538–1546.
- Wagner JA, Varga K, Ellis EF, Rzigalinski BA, Martin BR, Kunos G. Activation of peripheral CB1 cannabinoid receptors in haemorrhagic shock. *Nature*. 1997;390:518–521.
- Varga K, Wagner JA, Bridgen TD, Kunos G. Platelet- and macrophage-derived endogenous cannabinoids are involved in endotoxin-induced hypotension. *FASEB J*. 1998;12:1035–1044.
- Sugiura T, Kodaka T, Nakane S, Kishimoto S, Kondo S, Waku K. Detection of an endogenous cannabinoid molecule, 2-arachidonoylglycerol, and cannabinoid CB1 receptor mRNA in human vascular cells: is 2-arachidonoylglycerol a possible vasomodulator? *Biochem Biophys Res Commun*. 1998;243:838–843.
- Mechoulam R, Fride E, Ben-Shabat S, Meiri U, Horowitz M. Carbachol, an acetylcholine receptor agonist, enhances production in rat aorta of 2-arachidonoyl glycerol, a hypotensive endocannabinoid. *Eur J Pharmacol*. 1998;362:R1–R3.
- Zimmer A, Zimmer AM, Hohmann AG, Herkenham M, Bonner TI. Increased mortality, hypoactivity, and hypoalgesia in cannabinoid CB1 receptor knockout mice. *Proc Natl Acad Sci U S A*. 1999;96:5780–5785.
- Han L, Razdan RK. Total synthesis of 2-arachidonoylglycerol. *Tetrahedron Lett*. 1999;40:1631–1634.
- Rinaldi-Carmona M, Barth F, Millan J, Derocq J-M, Casellas P, Congy C, Oustric D, Sarrazin M, Bouaboula M, Calandra B, Portier M, Shire D, Breliere J-C, Le Fur G. SR144528, the first potent and selective antagonist of the CB2 cannabinoid receptor. *J Pharmacol Exp Ther*. 1998;284: 644–650.
- Goparaju SK, Ueda N, Yamaguchi H, Yamamoto S. Anandamide amidohydrolase reacting with 2-arachidonoylglycerol, another cannabinoid receptor ligand. *FEBS Lett*. 1998;422:69–73.
- Goparaju SK, Ueda N, Taniguchi K, Yamamoto S. Enzymes of porcine brain hydrolyzing 2-arachidonoylglycerol, an endogenous ligand of cannabinoid receptors. *Biochem Pharmacol*. 1999;57:417–423.
- Di Marzo V, Bisogno T, De Petrocellis L, Melck D, Orlando P, Wagner JA, Kunos G. Biosynthesis and inactivation of endocannabinoid 2-arachidonoylglycerol in circulating and tumoral macrophages. *Eur J Biochem*. 1999;264:258–267.
- Ben-Shabat S, Fride E, Sheskin T, Tamiri T, Rhee MH, Vogel Z, Bisogno T, De Petrocellis L, Di Marzo V, Mechoulam R. An entourage effect: inactive endogenous fatty acid glycerol esters enhance 2-arachidonoylglycerol cannabinoid activity. *Eur J Pharmacol*. 1998;353:23–31.